

Evidence for Extremely High Dust Polarization Efficiency in NGC 3184

Douglas C. Leonard

*Five College Astronomy Department, University of Massachusetts, Amherst, MA
01003-9305; leonard@nova.astro.umass.edu*

Alexei V. Filippenko, Ryan Chornock, and Weidong Li

*Department of Astronomy, University of California, Berkeley, California 94720-3411;
(alex, chornock, weidong)@astro.berkeley.edu*

ABSTRACT

Recent studies have found the Type II-plateau supernova (SN) 1999gi to be highly polarized ($p_{\max} = 5.8\%$, where $p_{\max}$ is the highest degree of polarization measured in the optical bandpass; Leonard & Filippenko 2001) and minimally reddened ($E[B-V] = 0.21 \pm 0.09$ mag; Leonard et al. 2002). From multiple lines of evidence, including the convincing fit of a “Serkowski” interstellar polarization (ISP) curve to the continuum polarization shape, we conclude that the bulk of the observed polarization is likely due to dust along the line of sight (l-o-s), and is not intrinsic to SN 1999gi. We present new spectropolarimetric observations of four distant Galactic stars close to the l-o-s to SN 1999gi (two are within 0.02°), and find that all are null to within 0.2%, effectively eliminating Galactic dust as the cause of the high polarization. The high ISP coupled with the low reddening implies an extraordinarily high polarization efficiency for the dust along this l-o-s in NGC 3184: $\text{ISP}/E[B-V] = 31^{+22}_{-9}\% \text{ mag}^{-1}$. This is inconsistent with the empirical Galactic limit ($\text{ISP}/E[B-V] < 9\% \text{ mag}^{-1}$), and represents the highest polarization efficiency yet confirmed for a single sight line in either the Milky Way or an external galaxy.

Subject headings: galaxies: individual (NGC 3184) — galaxies: ISM — supernovae: individual (SN 1999gi) — techniques: polarimetric

1. INTRODUCTION

It has long been understood that dust in the interstellar medium of the Milky Way (MW) is responsible for the linear polarization of starlight from intrinsically unpolarized

Galactic stars (see, e.g., Greenberg 1968, and references therein). The accepted model for the origin of the interstellar polarization (ISP) is linear dichroism (directional extinction), which results from aspherical dust grains aligned by some mechanism such that their optic axes have a preferred direction. By means of Mie theory computations (e.g., Greenberg 1968), the extinction efficiency factors for a specific grain size, shape, and refractive index may be calculated. Typically, the dust grains are modeled as dielectric cylinders with refractive index near 1.6, the value for silicates. The basic result is that an aspherical dust grain extinguishes slightly more light that is propagating with electric vector parallel to its long axis than it does light with electric vector perpendicular to its long axis. The resulting polarization direction is therefore perpendicular to the direction of the grain’s alignment.

The predicted degree of polarization produced by linear dichroism is strongly wavelength dependent, with maximum polarization occurring when the characteristic size of the grains (the cylinder’s radius) is of the same order as the wavelength of the light. Unlike the behavior of dust extinction, the expected spectral dependence of the ISP therefore rapidly decreases towards either longer *or shorter* wavelengths away from the peak. Such dust polarization models have been quite successful at reproducing the empirically derived “Serkowski law” ISP curve (Serkowski 1973; Wilking, Lebofsky, & Rieke 1982; Whittet et al. 1992) in the optical and near-infrared bandpasses (see, e.g., Mathis 1986).

Naturally, the same dust that polarizes the light is expected to redden it as well, a prediction observationally verified for MW stars. The degree of polarization produced for a given amount of extinction (or reddening) is referred to as the “polarization efficiency” of the intervening dust grains. The most efficient polarization medium conceivable is obtained by modeling the dust grains as infinite cylinders (length $\gg$ radius) with diameters comparable to the wavelength of the incident light, perfectly aligned with their long axes parallel to one another and perpendicular to the line-of-sight (l-o-s). For such a model, Mie calculations place a theoretical upper limit on the polarization efficiency of the grains due to directional extinction at visual wavelengths of (Whittet 1992, and references therein)

$$\frac{\text{ISP}}{E(B-V)} < R_V \times 13.82\% \text{ mag}^{-1}, \quad (1)$$

where ISP is the maximum degree of interstellar polarization measured in the optical band, $E(B-V)$ is the reddening, and R_V is the ratio of total to selective extinction (e.g., Savage & Mathis 1979). This upper limit on the polarization efficiency does not vary greatly with the refractive index of the grains considered, and is therefore rather insensitive to uncertainty in grain composition (Spitzer 1978, p. 175). Adopting the canonical value of $R_V = 3.1$ for MW dust leads to a maximum theoretical polarization efficiency along typical lines of sight in the MW of $\text{ISP}/E(B-V) < 43\% \text{ mag}^{-1}$. This may be compared with the empirically derived

upper bound (Serkowski, Mathewson, & Ford 1975) resulting from polarization studies of reddened Galactic stars of

$$\frac{\text{ISP}}{E(B-V)} < 9.0\% \text{ mag}^{-1}. \quad (2)$$

The fact that the observed polarization efficiency in the MW is more than a factor of 4 less than theory allows is generally taken as evidence that the alignment of dust grains is not total (or has multiple preferred orientations due to non-uniformity of the magnetic field along the l-o-s), and/or that the grains are only moderately elongated particles (rather than infinite cylinders) that may be irregularly shaped.

The empirical limit given by Equation (2) is found to hold for thousands of lines of sight to Galactic stars. To be sure, there are several stars in the comprehensive stellar polarization catalog by Heiles (2000) that exceed this limit (indeed, several polarized stars have claimed values of $E(B-V) \leq 0.0$ mag, formally creating an infinite polarization efficiency), but the colors and spectral types of the stars are not generally thought to be determined with sufficient accuracy to claim a clear violation of the inequality given by Equation (2).

Somewhat surprisingly, there have been very few measures of the polarization efficiency of dust in external galaxies. By far the best way to determine the polarization efficiency of dust in an external galaxy is to measure the ISP along the l-o-s to a discrete source with a well-determined reddening. Although this has been a very rare exercise (see Hough 1996 for a review of the four studies known at that time, and Rodrigues et al. 1997 for a subsequent study of the polarization and reddening of stars in the Small Magellanic Cloud), the results of the investigations that have been carried out are generally consistent with the upper limit given by Equation (2); while a few of the Large Magellanic Cloud stars studied by Clayton, Martin, & Thompson (1983) do exceed the limit, the discrepancy has been attributed to incorrect spectral classification or intrinsic colors (Hough 1996).

As part of a campaign to study the physical geometry of young supernovae (SNe; Leonard, Filippenko, & Matheson 2000b; Leonard et al. 2000a, 2001, 2002a), Leonard & Filippenko (2001; hereafter LF01) present a single epoch of optical spectropolarimetry of the bright, Type II-plateau supernova (SN) 1999gi in NGC 3184 during the late photospheric phase (107 days after discovery). They find an extraordinarily high degree of linear polarization, $p_{\text{max}} = 5.8\%$, where p_{max} is the highest level of polarization observed in the optical bandpass. If intrinsic to SN 1999gi, such polarization is without precedent for this type of event (see Wheeler 2000; Leonard et al. 2001; LF01). It implies an enormous departure from spherical symmetry, since even the most strongly aspherical theoretical models (axis ratio of 5-to-1, favorably viewed edge-on) of Höflich (1991) fail to produce polarizations of more than $\sim 5\%$. However, there is compelling evidence that much of the observed polarization is, in fact, due to interstellar dust. One clue pointing toward an interstellar origin is that the

polarization is not flat with wavelength, but rather has a broad, asymmetric peak that gently decreases on either side (Fig. 1). This is unlike the wavelength-independent polarization expected from an aspherical electron-scattering supernova atmosphere, but is very similar to the polarization observed for reddened Galactic stars (Serkowski 1973; Wilking et al. 1982; Whittet et al. 1992). Since the same dust that polarizes the SN light should also redden it, the natural expectation is for significant reddening. But, the Galactic foreground reddening being only $E(B-V) = 0.017$ mag (Schlegel, Finkbeiner, & Davis 1998), LF01 contend that nearly all of the dust along the l-o-s to SN 1999gi must reside in the host galaxy, NGC 3184.

Leonard et al. (2002c; hereafter L02) present optical spectra and photometry of SN 1999gi spanning the first six months after discovery as part of a study to derive the distance to SN 1999gi through the expanding photosphere method. Surprisingly, L02 conclude that SN 1999gi is only minimally reddened: Estimates resulting from five independent techniques are all consistent with a hard upper limit of $E(B-V) < 0.45$ mag established by comparing the early-time color of SN 1999gi with that of an infinitely hot blackbody, and yield a probable reddening of $E(B-V) = 0.21 \pm 0.09$ mag.

In this paper, we further investigate the nature of the observed polarization of SN 1999gi reported by LF01 in light of the recent evidence for low reddening found by L02. In § 2 we present new spectropolarimetric observations of four distant Galactic “probe” stars close to the l-o-s of SN 1999gi in order to test the assertion by LF01 that the polarization is not due to Galactic dust, and discuss the results in § 3.1. In § 3.2 we consider alternatives to ISP as the cause of the polarization, and introduce a new technique to quantify the component of SN polarization that lies in the direction perpendicular to the ISP direction in the Stokes parameter q - u plane. In § 3.3 we derive the polarization efficiency for dust in NGC 3184 along the l-o-s to SN 1999gi. We discuss the results in § 4 and summarize our conclusions in § 5.

2. Observations and Results

In order to investigate the contribution of Galactic ISP to the high polarization reported by LF01 for SN 1999gi, we obtained spectropolarimetry of four distant Galactic “probe” stars close to the l-o-s of SN 1999gi: BD +42°2113, HD 89572, and two “local standard stars” used by L02 to calibrate the photometry (labeled B and C in Fig. 2). The data were obtained on 2001 October 25 (BD +42°2113 and HD 89572) and 2002 January 8 (Stars B and C), using the Kast double spectrograph (Miller & Stone 1993) with polarimeter at the Cassegrain focus of the Shane 3-m telescope at Lick Observatory. The data were reduced according to the methods detailed by Leonard et al. (2001), and the results are given in Table 1.




Fig. 1.— Observed polarization (calculated as the rotated Stokes parameter; see Leonard et al. 2001) of SN 1999gi (*thin line*) on 2000 Mar. 25 (107 days after discovery; UT dates are used throughout this paper) compared with the total flux spectrum from the same date (*thick line*), arbitrarily scaled and offset for comparison of features. A Serkowski ISP curve characterized by $p_{\max} = 5.78\%$, $\theta = 154^\circ$, and $\lambda_{\max} = 5100 \text{ Å}$ is shown for comparison (*dashed line*; see § 3.2). Prominent features in the spectropolarimetry and total flux spectra are labeled.

Table 1. Probes of Galactic Dust Polarization Near the Line of Sight to SN 1999gi

Star	Spectral Type ^a	$m_V - A_V$ ^b (mag)	M_V ^c (mag)	Distance ^d (pc)	Angular Separation ^e (deg)	p_V (percent)	θ_V (deg)
BD +42°2113	A _m	8.83	0.6	$> 440 \pm 100$	0.52	0.07 ± 0.01	125 ± 3
HD 89572	A0	6.72	0.6	$> 170 \pm 40$	0.55	0.03 ± 0.01	98 ± 8
Star B	$\sim$ G6	15.41	5.5	$> 960 \pm 220$	0.02	0.19 ± 0.07	73 ± 9
Star C	$\sim$ G0	11.92	4.5	$> 300 \pm 70$	0.02	0.10 ± 0.02	111 ± 5

Note. — Stars B and C are the local standards labeled in Figure 2. The uncertainty in the polarization values is statistical only, and does not incorporate the 0.1 – 0.2% systematic uncertainty that may exist in polarimetry measurements made with the Lick 3 m polarimeter (see § 2). For low polarization values, the relatively large systematic uncertainty makes θ_V much more uncertain than the statistical error alone implies. Note that p_V and θ_V represent the debiased, flux-weighted averages of the polarization and polarization P.A., respectively, over the wavelength range 5050 – 5950 Å (see Leonard et al. 2001).

^aTaken from Simbad for BD +42°2113 and HD 89572, and estimated from colors and spectral features for Star B and Star C; see text for details.

^b $A_V = 0.05 \pm 0.01$ mag (Schlegel et al. 1998) adopted for all stars; m_V taken from Simbad for BD +42°2113 and HD 89572, and from L02 (Table 1) for Stars B and C.

^cFor luminosity class V dwarfs, from Binney & Merrifield (1998, p. 107). For BD +42°2113, the M_V for an A0 star is quoted, since this classification is most consistent with its observed $B - V$ color.

^dEstimated from spectroscopic parallax with the assumption that all stars are luminosity class V dwarfs; this assumption results in lower bounds on the inferred distances. The quoted uncertainty is derived by assuming a 0.04 mag uncertainty for $(m_V - A_V)$ and a 0.5 mag uncertainty for M_V (a rather liberal estimate, arising from the difficulty inherent in exact spectral classification).

^eBetween the probe star and SN 1999gi.

To determine the polarization angle (P.A.) offset between the half-wave plate and the sky coordinate system on 2001 October 25, we observed the polarized standard star BD +59°389 and set its V -band polarization position angle (i.e., θ_V , where the V subscript denotes the debiased, flux-weighted average over the wavelength range 5050 – 5950 Å; see Leonard et al. 2001) equal to 98.09°, the value cataloged by Schmidt, Elston, & Lupie (1992). To check the precision of the P.A. offset we also observed the polarized standard star HD 19820 on this night, and found its polarization angle to agree to within 0.2° of the value measured by Schmidt et al. (1992). The observed null standard, BD +32°3739 (Schmidt et al. 1992), was found to be null to within 0.1%.

To determine the P.A. offset on 2002 January 8, we observed HD 58624 and set θ_V equal to 26.9° (Mathewson & Ford 1970). To check the precision of the P.A. offset on this night we also observed the polarized standard stars HD 43384 and HD 245310. For HD 43384, θ_V was measured to be 170.8°, which agrees to within $\pm 0.3^\circ$ of the value cataloged by Mathewson & Ford (1970). However, the V -band polarization angle of HD 245310 was found to be 148.0°, which disagrees by 2° with the value ($\theta_V = 146^\circ$) reported by Hiltner (1956). While a 2° disagreement is not very alarming (and, in fact, is not that uncommon when comparing with older catalogs), it is also possible that a small amount of instrumental polarization existed on this night: the observed null standard, HD 57702, was measured to have $p_V = 0.17\%$, which is 0.13% higher than the value reported by Mathewson & Ford (1970).

The slight polarization for the null standard on 2002 January 8 could also be explained by systematic uncertainty. From repeated observations of a single standard star, Leonard et al. (2001) estimate the systematic uncertainty of the Lick polarimeter to be $\gtrsim 0.05\%$ in a Stokes parameter, or roughly 0.1% in the determination of p_V . In addition, Leonard et al. (2001) show that the uncertainty rises for objects with complex backgrounds, such as SNe in dusty galaxies or Galactic stars superposed on such backgrounds. Repeated observations of SN 1999em, in fact, show disagreements in a single Stokes parameter of up to 0.2%. Given these uncertainties, we shall consider any measured polarization that is $< 0.2\%$ to be effectively null. Fortunately, although uncertainty at this level certainly places a cautionary note on the interpretation of very low polarization measurements, it does not significantly hamper the interpretation of large polarization values, such as that measured for SN 1999gi.

In order to serve as good probes of interstellar dust polarization, it is clearly necessary to observe stars that lie significantly above (or below) the Galactic plane and close to the l-o-s of the SN. Applying the criterion of Tran (1995) that a good MW “probe” star should be more than 150 pc from the Galactic plane, the Galactic latitude of SN 1999gi ($b \approx 55^\circ$) necessitates using stars that are at least 183 pc away, a criterion that all but one of our probe stars (HD 89572, $d = 170 \pm 40$ pc) definitely meets (see Table 1). We estimated the




Fig. 2.— *B*-band image of NGC 3184 taken on 1999 December 17 with the Katzman Automatic Imaging Telescope (KAIT; Filippenko et al. 2001), with the four local standards used by L02 marked. Spectropolarimetry of Stars B and C (both within 0.02° of the l-o-s to SN 1999gi and distant enough to fully sample Galactic dust) establishes that both are effectively null ($p_V < 0.2\%$), essentially eliminating Galactic dust as the cause of the high polarization found by LF01 for SN 1999gi. The line segment through SN 1999gi (*SN*) at P.A. = 154° indicates the polarization angle of the SN measured by LF01.

probe star distances through spectroscopic parallax, with the spectral types and apparent magnitudes for BD +42°2113 and HD 89572 taken from the Simbad Astronomical Database.¹ To determine the spectral types of the two “anonymous” local standard stars we compared their observed colors, corrected for a Galactic reddening of $E(B-V) = 0.017$ mag (Schlegel et al. 1998; $[B-V] = 0.69$ and 0.57 mag for Stars B and C, respectively), with the values tabulated by Binney & Merrifield (1998, p. 107) for luminosity class V dwarfs. (Note that if either of the probe stars is in fact more intrinsically luminous than a class V dwarf, it would serve to increase the inferred distance, which only enhances their efficacy as probe stars). We also verified this approximate classification through examination of their spectra (Fig. 3). With the possible exception of HD 89572, then, all of the probe stars lie considerably farther away than the Tran (1995) criterion requires, and are sufficiently close to the l-o-s to SN 1999gi to serve as good probes of the Galactic ISP.

3. Analysis

3.1. Galactic Polarization

From Table 1 we see that all of the probe stars are effectively null, confirming the suspicion of LF01 that the cause of the high polarization is not local to the MW, but rather resides in the host galaxy, NGC 3184. This is also consistent with the low Galactic reddening ($E[B-V] = 0.017$ mag) reported by the dust maps of Schlegel et al. (1998). Since Stars B and C are located on either side of SN 1999gi (Fig. 2) and are only $\sim 0.04^\circ$ away from each other, this constrains the possible radius of an undetected polarizing spherical Galactic dust blob (or one dimension of a filamentary structure) that covers only SN 1999gi and neither probe star to values less than ~ 0.13 pc, assuming that the undetected blob resides within 150 pc of the Galactic plane. While such tiny concentrations of dust may be possible (e.g., the outer edge of a Bok globule, or a thin filamentary dust structure), the chance for such an alignment is certainly quite remote. We therefore feel confident in asserting that Galactic dust is not responsible for the polarization of SN 1999gi.

3.2. Alternatives to ISP

Before concluding that the high polarization of SN 1999gi is due to ISP from dust in NGC 3184, however, we consider three alternatives.

¹<http://simbad.u-strasbg.fr/Simbad>




Fig. 3.— Optical flux spectra of the two local dust probe stars discussed in the text and labeled in Figure 2, with two additional spectra shown for comparison of features. All spectra have been normalized so that the $H\alpha$ line has the same depth in each spectrum. The strength of the neutral metal lines (e.g., the Ca I lines indicated) in stars of type K2V (HD 3765) and later effectively restricts the classification of the two probe stars to earlier types. This is consistent with the color-based classification of the two stars of G0 (Star C) and G6 (Star B). Note that the resolution of the two comparison spectra ($\sim 2.5 \text{ \AA}$) is somewhat higher than that of the local probe stars ($\sim 6 \text{ \AA}$). The comparison spectra were taken using the Keck II 10-m telescope with the Low Resolution Imaging Spectrometer (Oke et al. 1995) as part of another program; see, e.g., Filippenko et al. (1999).

1. *The polarization is due to newly formed dust in the expanding SN ejecta.* It is now generally accepted that dust formation is a natural process in evolved SN atmospheres (e.g., Gerardy et al. 2000). However, there are several objections to this proposal for SN 1999gi at the epoch studied here, about 111 days after the explosion (L02). First, the temperature of the ejecta at the observed phase, including the inner regions in which the dust is thought to ultimately form (e.g., Wooden 1997; Clayton, Deneault, & Meyer 2001), is greatly in excess of the ~ 2000 K evaporation temperature of iron-rich grains (Wooden 1997), making it a very hostile environment for dust survival (Rank et al. 1988; Dwek 1991; Wooden et al. 1993). Indeed, the detailed study of SN 1987A by Wooden (1997) shows that the earliest possible epoch for dust to have formed in that event is about 350 days after explosion. In addition, the spectral and photometric evolution of SN 1999gi do not exhibit any of the telltale optical signatures of dust formation in the time leading up to the spectropolarimetric observation, such as an accompanying drop in optical luminosity or the onset of asymmetric line profiles (e.g., SN 1998S; Leonard et al. 2000). Dust formation in the ejecta of SN 1999gi can therefore be confidently eliminated as a possibility at this comparatively early evolutionary phase.

2. *The polarization is produced by dust reflection by one or more off-center dust blobs in NGC 3184 near to the l-o-s of SN 1999gi.* Reflection by dust is highly polarizing, and certainly could produce the average level of polarization seen in SN 1999gi. However, reproducing the observed Serkowski-law dependence, while surely possible given all conceivable dust-blob orientations and distances, would certainly require some fine-tuning of the model since simple reflection (i.e., single scattering) off grains of typical size produces polarization that increases at blue wavelengths due to the greater scattering efficiency there (e.g., Miller & Goodrich 1990; Miller, Goodrich, & Mathews 1991; see Kartje 1995 for a discussion of how multiple scatters could alter this). While computing a detailed light-echo model (e.g., Wang & Wheeler 1996) is beyond the scope of this study, we consider the possibility of dust reflection mimicking a Serkowski-law ISP curve to be quite remote.

3. *The observed polarization is intrinsic to SN 1999gi itself.* It is clear that SN 1999gi has some intrinsic polarization. The distinct spectropolarimetric features seen in Figure 1 correspond directly to strong features in the total flux spectrum; such sharp polarization modulations are not characteristic of ISP and must belong to the SN. There are, however, several reasons to believe that the intrinsic SN polarization component is relatively minor compared with the ISP.

To start with, the strength of the line features relative to the continuum polarization is quite small. The basic paradigm used to interpret polarization changes across P-Cygni line features is that polarization peaks are naturally associated with absorption minima due to

selective blocking of forward-scattered (and hence less polarized) light in P-Cygni absorption troughs and polarization minima are associated with emission peaks due to the dilution of polarized continuum light with unpolarized line emission (e.g., McCall 1984; Jeffery 1991a; LF01; Leonard et al. 2002a). In fact, the polarization level in the emission components of strong lines (e.g., $H\alpha$, Ca II infrared triplet) has been used in several previous studies to define the ISP level (e.g., Trammell, Hines, & Wheeler 1993; Leonard et al. 2001, 2002a; Kawabata et al. 2002). For SN 1999gi, the lack of any strong polarization change across the $H\alpha$ emission component suggests that the ISP level is not significantly different from the observed polarization. In the absorption troughs, we see modest modulations at the locations of Ba II $\lambda 4554$, $H\beta$, Na I D, and possibly Fe II $\lambda 5169$ as well (Fig. 1). Curiously, the polarization changes in the troughs are observed to be *decreases* rather than the expected increases; we shall discuss the implications of this unusual behavior later in this section. For now, we note that LF01 find the strength of the modulations to be consistent with other SNe II-P having low intrinsic continuum polarization, and conclude that they are consistent with an intrinsic continuum polarization as low as $p_{\text{SN}} \approx 0.3\%$ for SN 1999gi.

Another argument against high intrinsic SN polarization is that the continuum polarization shape is not flat with wavelength, as would be expected for an aspherical electron-scattering atmosphere, but rather approximates a Serkowski law ISP curve. In Figure 1 we show one possible Serkowski law fit to the data, determined by finding the best agreement (lowest χ^2) between a Serkowski law ISP curve (Whittet et al. 1992) and the observed spectropolarimetry in wavelength regions not obviously affected by line-trough features (we chose to fit the regions 4600–4810 Å, 5000–5100 Å, 5230–5670 Å, and 6070–6460 Å). The derived Serkowski law ISP curve shown in Figure 1 is certainly not a unique solution: the spectral regions that were fit were chosen somewhat arbitrarily (indeed, the “best” fit presented in Fig. 1 is slightly different from that derived by LF01, due largely to the different spectral regions that were fit) and *some* intrinsic SN polarization definitely exists, which makes the true ISP level and spectral shape slightly different from that to which we fit. Nonetheless, the good qualitative agreement between the overall shape of the observed polarization and a Serkowski law ISP curve, and its incompatibility with wavelength-independent polarization, is quite suggestive that ISP, and not intrinsic SN polarization, dominates the observed polarization.

The constancy of the polarization angle with wavelength across the optical spectrum (θ never strays by more than 2° from 154° ; see Fig. 8 of LF01) also casts doubt on large intrinsic SN polarization since, unless the SN polarization P.A. is either parallel or perpendicular² to the ISP P.A., a somewhat unlikely chance occurrence, significant variation

²Recall that $\theta \equiv 1/2 \arctan(u/q)$, so that a P.A. difference of 90° corresponds to a 180° orientation




Fig. 4.— Polarization data for SN 1999gi obtained on 2000 March 25, where each point represents a bin of $100 \text{ \AA}$ in each of the Stokes parameters. The best-fitting line, determined by a uniform-weight least-squares fit, is drawn through the points, along with the line perpendicular to it that intersects the origin. The distance between the point of intersection of these two lines and the origin reveals the component of wavelength-independent SN continuum polarization in the direction perpendicular to the best-fitting line in the q - u plane, $p_{\perp} = 0.29 \pm 0.43\%$. Note that the statistical error of each point is insufficient to explain the scatter about the best-fitting line, which is likely produced by the effect of low-contrast line features in the spectropolarimetry (Fig. 1).

in the polarization angle with wavelength would be observed. To quantify this assertion, we consider the representation of polarization data in the q - u plane (Fig. 4). Unlike the wavelength-independent nature of intrinsic SN continuum polarization, which will produce a spherical cluster of points in the q - u plane, a Serkowski law wavelength-dependence of ISP should produce points that are spread out along a given P.A.; due to noise, systematic uncertainty, and the effect of line features, the points will not all lie exactly on a line, but will instead have some scatter about it. In the case of pure, undiluted ISP from a single source, we would expect the best-fitting line to the points to intersect the origin in the q - u coordinate system, since this is the definition of having a single P.A. If another source possessing a significant polarization component at $\pm 45^\circ$ to the ISP P.A. is contributing, then the best-fitting line will not intersect the origin. Under the simple paradigm of a single ISP P.A. and wavelength-independent SN continuum polarization, in fact, the amount by which the best-fitting line misses the origin reveals the component of SN polarization oriented at $\pm 45^\circ$ to the ISP P.A. (i.e., $p_\perp \equiv \sqrt{q_o^2 + u_o^2}$, where q_o and u_o are the points of intersection in the q - u plane between the line defined by the polarization data and the line perpendicular to it that passes through the origin).

Figure 4 shows the polarization data for SN 1999gi, where each of the 25 points represents a 100 Å bin in the q and u Stokes parameters. The best-fitting line, determined by a uniform-weight least-squares fit, is seen to miss the origin by only $p_\perp = 0.29\%$. The statistical uncertainty of the individual points are clearly insufficient to explain the scatter about the best-fitting line, which is likely produced by the effect of low-contrast line features in the spectropolarimetry (Fig. 1). Therefore, to determine the uncertainty in the value of $p_\perp$, we employed the technique of bootstrap resampling (Press et al. 1992), in which new data sets of 25 points were created by randomly drawing data values from the original 25 data points with repetition of values allowed. We then computed $p_\perp$ for 1000 data sets resampled in this way and found the 1σ scatter of the results to be 0.43%, which we adopt as the 1σ uncertainty on the value of $p_\perp$. This convincingly demonstrates that significant (i.e., more than $\sim 1.5\%$) wavelength-independent intrinsic SN polarization must have a polarization angle either coincident with, or perpendicular to, the ISP direction.

With this in mind, we now return to the curious observation that the observed polarization of SN 1999gi *decreases* in the absorption troughs of strong lines in the total flux spectrum (see Fig. 1), contrary to what is expected theoretically and also to what has been observed in the spectropolarimetry of other core-collapse SNe (e.g., SN 1987A [Jeffery et al. 1991b, and references therein]; SN 1993J [Tran et al. 1997; Höflich et al. 1996]; SN 1997ds [LF01];

difference in the q - u plane.

SN 1999em [Leonard et al. 2001]; SN 2002ap [Leonard et al. 2002a; Kawabata et al. 2002]). The explanation for such atypical behavior is straightforward, however, if ISP dominates the observed polarization and the SN polarization has a P.A. that is oriented perpendicular to the P.A. of the ISP (as we suspect may be the case from the earlier q - u plane analysis). That is, if the intrinsic SN polarization has a P.A. that lies along the best-fitting line in Figure 4, but in the upper left quadrant of the q - u plane (i.e., at P.A. $\approx 64^\circ$, perpendicular to the ISP P.A. of $\sim 154^\circ$), then *intrinsic* SN polarization level increases would become polarization decreases in the *observed* spectropolarimetry due to the ISP. The unusual polarization decreases in the line troughs may therefore be only an artifact of a much stronger contribution from the ISP to the overall polarization level.

A final argument against large intrinsic SN polarization is the fact that no SN II-P has ever been found to be intrinsically polarized at more than 1.5% (Wheeler 2000; LF01; Leonard et al. 2001). Indeed, in the detailed multi-epoch spectropolarimetric study by Leonard et al. (2001) of SN 1999em, shown by L02 to be a spectral and photometric twin of SN 1999gi, the polarization does not rise above $p_V = 0.5\%$ in the epochs studied (days 7, 40, 49, and 159 after discovery).

The preceding discussion therefore presents a choice: either SN 1999gi has the highest intrinsic polarization (by far) of any core-collapse SN yet observed, a continuum polarization shape that is unlike the wavelength-independent polarization expected from, and previously observed in, SNe II, and line polarization features unlike those observed in any previous SN II (i.e., polarization decreases in absorption troughs), or there is a large ISP component. The simplest explanation, and one that also restores normalcy to all of the peculiarities of the observed polarization, is that SN 1999gi possesses only minimal intrinsic polarization, and that a large ISP dominates. From the analysis of the line polarization and continuum shape, as well as the historical evidence against SNe II-P possessing high intrinsic polarization, we conclude that SN 1999gi is very likely to be intrinsically polarized in the range $0.3 \leq p_{\text{SN}} \leq 1.5\%$. We also note that if the majority of the intrinsic SN polarization is indeed oriented perpendicular to the P.A. of the ISP as we suspect, then the true ISP level is *even greater* than the level observed, due to the vector nature of polarization addition.

3.3. Polarization Efficiency of Dust in NGC 3184

The most viable production mechanism for the bulk of the polarization observed in SN 1999gi is therefore directional extinction by interstellar dust grains along the l-o-s in NGC 3184. Assuming the wavelength of maximum ISP to be $\lambda_{\text{max}} \approx 5100 \text{ \AA}$ (§ 3.2; Fig. 1) and adopting the empirical correlation between the wavelength of maximum polarization

and R_V found by Whittet & van Breda (1978), we derive $R_V = 2.9$ for the dust polarizing the light of SN 1999gi (cf. LF01). The theoretical upper bound on the dust polarization efficiency from Equation (1) is therefore

$$\frac{\text{ISP}}{E(B-V)} < 40\% \text{ mag}^{-1}. \quad (3)$$

If the dust has a polarization efficiency similar to typical MW dust, of course, then the empirical upper limit given by Equation (2) is expected to hold.

From the limits on the intrinsic SN polarization established in § 3.2 as well as the null results for the Galactic probe stars, we conclude that the polarization contributed by dust in NGC 3184 likely lies in the range $4.3\% \leq \text{ISP} \leq 7.3\%$. (We note that the nature of the line trough polarization discussed in § 3.2 actually favors restricting the ISP even further, to values greater than 5.8%; for the purpose of calculating the range of *possible* polarization efficiency, however, it seems prudent to adopt the most liberal limits.) When combined with the reddening of $E(B-V) = 0.21 \pm 0.09$ mag (L02), we derive a polarization efficiency³ for the dust in NGC 3184 of $31^{+22}_{-9}\% \text{ mag}^{-1}$. Figure 5 displays the range of possible values for the ISP and reddening for the dust in NGC 3184 along the l-o-s to SN 1999gi, with the empirical and theoretical limits prescribed by Equations (2) and (3), respectively, indicated as well. For lack of a clear physical motivation for favoring a specific ISP value within the allowed range, we nominally place it at $\text{ISP} = 5.8\%$, the observed polarization value.

The range of possible ISP and reddening values shown in Figure 5 is inconsistent with the maximum observed Galactic polarization efficiency. Our nominal value is three times the

³Due to the non-Gaussian nature of the ISP probability distribution (i.e., it is flat between the end points), the quoted values for the polarization efficiency and uncertainty were derived through a Monte Carlo simulation. We first randomly selected 10,000 values from a Gaussian-weighted distribution for the reddening ($E[B-V] = 0.21 \pm 0.09$ mag) and a flat probability distribution for the ISP ($4.3 \leq \text{ISP} \leq 7.3$). From physical considerations (i.e., dust is never observed to *deredden* light) we did not allow negative reddening. So that the distribution of reddening values would remain symmetric about $E(B-V) = 0.21$ mag, we also did not allow reddenings above 0.42 mag. This latter restriction is quite justifiable as well from the upper limit of $E(B-V) < 0.45$ mag derived by L02. We therefore replaced any reddenings that were not within the range $0.0 < E(B-V) < 0.42$ mag with ones drawn randomly from the Gaussian-weighted reddening distribution that do fall within the allowed range. We then computed the polarization efficiency for the 10,000 sets of values. The resulting probability distribution is not normally distributed, but rather has positive skewness and a few extraordinarily large values resulting from very small (but allowable) reddenings. These outlier points bias both the mean and the standard deviation to large, non-robust values (i.e., they cannot be reproduced from one simulation to the next). Our quoted result is therefore the *median* of the probability distribution, and the lower and upper values of the uncertainty mark the locations of the 16th and 84th percentile points, respectively, of the simulated set of data (i.e., the 68% confidence bounds).




Fig. 5.— The ISP and reddening along the l-o-s to SN 1999gi in NGC 3184, with the observed polarization and best reddening estimate indicated by the *filled circle*. The 1σ reddening uncertainty derived by L02 is indicated (*short error bars*), as are the lower and upper limits for both the reddening and ISP (*long error bars*). Also shown is the empirical limit on the polarization efficiency of MW dust (*dotted line*; Serkowski et al. 1975), as well as the theoretical upper limit for dust grains consisting of completely aligned infinite dielectric cylinders (*solid line*; Whittet 1992).

empirical limit, and the upper end of the range extends beyond the theoretical maximum given by Equation (3). Even the limiting values of $E(B-V) = 0.45$ mag and $ISP = 4.3\%$ produce a polarization efficiency above (but just barely) the Galactic limit, and these limiting values are thought to be quite unlikely. Therefore, unless the estimate of the reddening or the allowed range of the ISP are grossly in error, this measurement is the *highest polarization efficiency yet confirmed for a single sight line in either the MW or an external galaxy*.

4. Discussion

4.1. Galactic ISP Models

What does the high polarization efficiency tell us about the dust in NGC 3184? Does it suggest an unusually high degree of grain alignment? An unusually strong magnetic field? Grain shapes that actually resemble infinite cylinders? To gain insight we briefly discuss the various grain-alignment mechanisms that have been proposed for the MW over the years.

First, it is clear that the Galactic magnetic field plays a leading role in grain alignment, since the optical polarization position angles in the MW of the most distant stars (e.g., Mathewson & Ford 1970) are generally found to parallel the Galactic magnetic field direction inferred from other independent studies, largely at radio wavelengths (see, e.g., Heiles 1996 for a detailed comparison among the magnetic field directions derived from starlight polarization, pulsar rotation measure data, and synchrotron emission). Similar results have also recently been obtained using optical and infrared polarimetry of the *integrated* starlight of a limited number of external galaxies as well, such as NGC 4565, a normal spiral, in which the radio continuum (Sukumar & Allen 1991), optical polarimetry (Scarrott, Rolph, & Semple 1990), and infrared polarimetry (Jones 1989) are all consistent with a magnetic field lying in the plane of the galaxy.

The classic mechanism by Davis & Greenstein (1951) achieves alignment through paramagnetic relaxation of the thermally rotating grains in the presence of an external magnetic field. This mechanism, however, predicts a strong correlation between the degree of alignment and magnetic field strength (e.g., Aannestad & Purcell 1973; Johnson 1982), which is not observed (see, e.g., Heiles 1987 and references therein). For this and other reasons, the classic Davis-Greenstein mechanism is no longer seen as the most likely candidate for alignment, although a modification of the basic idea is still favored (e.g., Jones 1996 and references therein).

In recent years, the trend has been toward deemphasizing the role that magnetic field *strength* plays in grain alignment and instead focusing on its *direction*. In fact, a very suc-

successful recent model by Jones, Klebe, & Dickey (1992) posits no correlation of alignment with magnetic field strength at all (assuming, of course, that some nonzero field exists). Rather, the Jones et al. (1992) model assumes *complete alignment* of the large aspherical component of the grain population at all times (i.e., independent of the magnetic field strength), but with equal mixing of random and constant directional components to the magnetic field. That is, the model envisages a uniform Galactic magnetic field pervading an interstellar medium that itself is made up of individual cells containing superposed, randomly oriented magnetic field vectors with a strength comparable to that of the Galactic field. The random components have the effect of causing the polarization angle of the net field to vary from cell to cell, thereby reducing the overall polarization efficiency of the ISM.

The Jones et al. (1992) model successfully explains the correlation of polarization strength with extinction at $2.2\ \mu\text{m}$ (i.e., the K band), where it is possible to study stars that suffer over 100 mag of visual extinction. In addition to favoring equipartition between the random and constant components of the magnetic field, a prediction that is also in good agreement with radio observations (e.g., Heiles 1987, 1996), this model finds the extent of the cells to be dictated not by size, but, rather, by extinction. Specifically, the best agreement with the observations results when the random magnetic field component decorrelates over a path length of $A_V \approx 1$ mag, a result found to be valid for both diffuse gas and dense clouds. One implication of this is that stars with $A_V \lesssim 1$ mag should have higher polarization efficiency, on average, than those with greater extinction. High polarization efficiency, therefore, should prefer the very low-extinction regime, where the effect of only a single cell polarizes the light.

Within the context of the Jones et al. (1992) model, then, we might be able to explain the high polarization efficiency, the surprising polarization angle (i.e., that it is not aligned with the spiral arm; see, e.g., Scarrott et al. 1990, 1991), and the nearly constant wavelength dependence of the polarization angle if the bulk of the polarization of SN 1999gi is produced by a single dust cell of highly aligned grains with $l \gg r$ whose orientation is determined primarily by a local, randomly oriented magnetic field. Even with these specifications, however, it is possible that an unusually regular magnetic field and/or different alignment mechanism may be required to explain such high polarization efficiency.

4.2. The Polarization Efficiency of Galactic Dust

Although the low-extinction regime is only sparsely sampled by the data of Jones et al. (1992; only 3 of the 99 stars studied have $A_V \lesssim 1$ mag, and none is found to violate the empirical Galactic limit set by Equation [2]), it is instructive to review the low-extinction

polarization data that *are* available for Galactic stars, to see if the extraordinary polarization efficiency indicated by SN 1999gi is unique.

The most complete collection of stellar polarization data is the agglomeration of stellar polarization catalogs by Heiles (2000), which contains polarization and, where available, reddening data for 9286 Galactic stars; in nearly all cases, the quoted stellar reddening was derived in the original catalog by comparing a star’s intrinsic color (estimated from its spectral type) with the observed color. An investigation of the low-extinction regime of the Heiles (2000) catalog is given by Fosalba et al. (2002), in which only stars with listed degrees and angles of polarization, as well as positive extinctions, are studied (several stars in the Heiles (2000) catalog have negative reddenings, since these values were reported by the original catalogs from which Heiles’ data were drawn); this subsample constitutes $\sim 60\%$ of the stars in the Heiles (2000) catalog. When looking at the average polarization properties of the stars, Fosalba et al. (2002) indeed find evidence for increasing average polarization efficiency with decreasing reddening in the reddening range $E(B-V) < 1.0$ mag: $P_{\text{avg}}/P_{\text{max}} = 0.39E(B-V)^{-0.2}$, where P_{avg} is the average value of the polarization and P_{max} is the empirical limit given in Equation (2) for each reddening. Although a few dozen stars are found to exceed the Serkowski et al. (1975) limit, Fosalba et al. (2002) pass over these violations without comment, perhaps due to the inherent uncertainties in the techniques used to gauge stellar reddenings (e.g., Teerikorpi 1990), a difficulty that has added caution in the interpretation of polarization efficiency deduced by other researchers as well (e.g., Berdyugin, S      , & Teerikorpi 1995). An additional challenge when interpreting polarization data in the very low-extinction regime is the bias toward high polarizations that results from the positive-definite definition of polarization (i.e., $p = \sqrt{q^2 + u^2}$) as well as the positive skewness exhibited by its probability distribution (e.g., Miller et al. 1988; see Leonard et al. 2001 for a thorough discussion of these issues).

To more securely investigate possible violations of Equation (2) at low reddenings, the recent dust maps of Schlegel et al. (1998) offer a useful alternative to extinctions estimated by stellar color excess. First, they provide reddenings along Galactic lines-of-sight accurate to $\sim 16\%$. Second, they represent the reddening produced by the *total* Galactic dust column along a particular l-o-s, so that the value serves as a natural upper bound on the reddening to any Galactic star. Using the Schlegel et al. (1998) reddening values will therefore result in *lower bounds* on the derived polarization efficiencies of Galactic stars. Thus, any star whose polarization efficiency is found to significantly violate Equation (2) when calculated in this way can be flagged as a potential exception to the Serkowski et al. (1975) limit. One point that must be kept in mind, however, is that the Schlegel et al. (1998) maps have a resolution of $\sim 6'.1$ which, though capable of resolving quite small-scale features, certainly leaves open the possibility of unresolved dust filaments. Nonetheless, the exercise is worthwhile as a

preliminary investigation.

Figure 6 shows the results of applying the Schlegel et al. (1998) reddenings to the polarization data of Heiles et al. (2000) in the low-extinction regime, and several stars clearly do violate the Serkowski et al. (1975) limit. Although we did not conduct a detailed investigation of all of these stars, we did attempt to further investigate the most egregious violators, for which the observed polarization exceeds the empirical limit by more than 2% (stars 3200, 3340, 3355, 3366, 3530, 3543, and 5997 in the Heiles 2000 list; note that the dust along the l-o-s to SN 1999gi likely violates the empirical limit by $\gtrsim 2.5\%$). Unfortunately, the source of the data for all of these exceptional stars is an unpublished catalog by A. Goodman (see Heiles 2000 and references therein) for which only positions and measured polarizations, and no uncertainties or reddenings, are given. This catalog contains stars primarily in, near, or behind star-forming molecular clouds, where the density of H_2 molecules exceeds $10^2 - 10^3 \text{ cm}^{-3}$; such regions are noticeable as dark patches of extinction on optical photographs and are therefore often referred to as “dark clouds”. From the given positions we determined that all but the last star (Star 5997) lie in the vicinity of the dark cloud L1755, a star-forming region in Ophiuchus, for which additional information for two of the stars, numbers 3530 and 3543, are given by Goodman et al. (1990) as $p = 5.87 \pm 0.12\%$ and $p = 6.02 \pm 0.13\%$, respectively. The polarization values and stated uncertainties certainly place these stars well above the empirical limit for their Schlegel et al. (1998) reddenings (0.38 mag and 0.40 mag for Star 3530 and Star 3543, respectively).⁴ At an estimated distance of ~ 160 pc to the L1755 cloud (Bertiau 1958) the dust maps of Schlegel et al. (1998) are capable of resolving structures of ~ 0.3 pc in size, which is sufficient to resolve the L1755 cloud itself, estimated to have a physical size of $\sim 6 \text{ pc} \times 0.7 \text{ pc}$ (Goodman et al. 1990). Certainly, though, it is conceivable that unresolved substructure exists in such an environment, and it is unfortunate that no specific extinction information exists for these stars.

Although the lack of specific reddening estimates precludes a definitive statement, the possibility certainly exists that violations of the empirical limit given by Equation (2) do occur in the MW. Additional studies of the low-reddening regime are needed before more can be said about Galactic violations of Equation (2), however. Until such studies are carried out, though, our measurement of the polarization efficiency of dust in NGC 3184, $\text{ISP}/E(B-V) = 31_{-9}^{+22}\% \text{ mag}^{-1}$, serves as the highest polarization efficiency yet confirmed for a single sight line in either the MW or an external galaxy.

⁴Note that the stars studied by Goodman et al. (1990) were specifically chosen to be coincident, in projection (i.e., they are most likely background stars), with the periphery of the extinction associated with the dark clouds. Such a selection criterion should naturally yield a sample with low inferred reddening.




Fig. 6.— Polarization of Galactic stars from the agglomeration catalog of Heiles (2000) plotted against the Galactic reddening given by the dust maps of Schlegel et al. (1998). Since the Schlegel et al. (1998) reddenings probe the entire Galactic dust column whereas the stars do not, the plotted reddenings represent upper bounds on the possible reddenings of these stars. The observed polarization and derived reddening of SN 1999gi is indicated by a *solid triangle*.

5. Conclusions

The single spectropolarimetric epoch obtained for the Type II-P SN 1999gi by LF01 indicates a very high observed polarization ($p_{\max} = 5.8\%$), with a shape that is well characterized by a Serkowski law ISP curve. The null result obtained by spectropolarimetric observations of four Galactic stars near to the l-o-s to SN 1999gi and sufficiently distant to fully sample Galactic dust convincingly eliminates Galactic ISP as a significant contributor to the observed polarization. It is also difficult to reconcile intrinsic SN polarization, newly formed dust in the ejecta, or dust reflection by off-axis dust blobs with the observed polarization characteristics. We therefore conclude that ISP produced by dust in NGC 3184 is largely responsible for the observed polarization, and that it lies in the range $4.3\% \leq \text{ISP} \leq 7.3\%$. Since L02 find SN 1999gi to be only minimally reddened, $E(B-V) = 0.21 \pm 0.09$ mag (with all reddening estimates supporting a firm upper limit of $E[B-V] < 0.45$ mag, which was derived through the comparison between the early-time continuum shape with that of an infinitely hot blackbody), we derive a polarization efficiency for the dust along this l-o-s in NGC 3184 of $31_{-9}^{+22}\% \text{ mag}^{-1}$. Even by assuming the limiting reddening value of $E(B-V) = 0.45$ mag and the lower ISP limit of $\text{ISP} = 4.3\%$, the derived polarization efficiency does not conform to the empirical Galactic limit of $\text{ISP}/E[B-V] < 9\% \text{ mag}^{-1}$, and these limiting values are thought to be extremely unlikely. Therefore, unless the estimates for the ranges of either the reddening or the ISP are grossly in error, this measurement is the highest polarization efficiency yet confirmed for a single sight line in either the MW or an external galaxy.

Although the polarization efficiency for dust along the l-o-s to SN 1999gi is inconsistent with the empirically derived maximum observed Galactic polarization efficiency, it can be accommodated by the maximum theoretical efficiency derived from Mie scattering considerations ($\text{ISP}/E[B-V] < 40\% \text{ mag}^{-1}$). In addition, the dust grain polarization model of Jones et al. (1992) predicts high polarization efficiency for lines of sight to sources with low reddening ($A_V < 1$ mag). From an examination of the Galactic stellar polarization agglomeration catalog of Heiles (2000), we also find some evidence for unusually high polarization efficiency for dust in dark clouds, although the current lack of specific extinction information for these regions precludes definitive conclusions. Since the low-reddening regime has been only sparsely sampled by previous studies, increased scrutiny of the polarimetry of discrete Galactic and extragalactic sources with low reddening certainly seems warranted to test whether the polarization efficiency inferred for dust along the l-o-s to SN 1999gi in NGC 3184 is indeed common in the low-reddening regime.

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France, and was supported in part by NASA through the American Astronomical Society’s

Small Research Grant Program. Our work was also funded by NASA grants GO-8648, GO-9114, and GO-9155 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS 5-26555. Additional funding was provided to A. V. F. by NASA/Chandra grant GO-0-1001C, by NSF grant AST-9987438, by the Guggenheim Foundation, and by the Sylvia and Jim Katzman Foundation. KAIT was made possible by generous donations from Sun Microsystems, Inc., the Hewlett-Packard Company, Auto-Scope Corporation, Lick Observatory, the National Science Foundation, the University of California, and the Katzman Foundation.

REFERENCES

- Aannestad, P. A., & Purcell, E. M. 1973, *ARA&A*, 11, 309
- Berdyugin, A., Snåre, M.-O., & Teerikorpi, P. 1995, *A&A*, 294, 568
- Bertiau, F. C. 1958, *ApJ*, 128, 533
- Binney, J., & Merrifield, M. 1998, *Galactic Astronomy* (Princeton: Princeton University Press)
- Clayton, D. D., Deneault, E. A.-N., & Meyer, B. S. 2001, *ApJ*, 562, 480
- Clayton, G. C., Martin, P. G., & Thompson, I. 1983, *ApJ*, 265, 194
- Davis, L., & Greenstein, J. L. 1951, *ApJ*, 114, 206
- Dwek, E. 1991, in *Supernovae*, ed. S. E. Woosley (New York: Springer-Verlag), 54
- Filippenko, A. V., Leonard, D. C., Matheson, T., Li, W., Moran, E. C., & Riess, A. G. 1999, *PASP*, 111, 969
- Filippenko, A. V., Li, W. D., Treffers, R. R., & Modjaz, M. 2001, in *Small Telescope Astronomy on Global Scales*, ed. W. P. Chen, C. Lemme, & B. Paczyński (San Francisco: ASP, Conf. Ser. Vol. 246), 121
- Fosalba, P., Lazarian, A., Prunet, S., & Tauber, J. A. 2002, *ApJ*, 564, 762
- Gerardy, C. L., Fesen, R. A., Höflich, P., & Wheeler, J. C. 2000, *AJ*, 119, 2968
- Greenberg, J. M. 1968 in *Nebulae and Interstellar Matter*, ed. B. M. Middlehurst & L. H. Aller (Chicago: University of Chicago Press), 221

- Heiles, C. 1987, in *Interstellar Processes*, ed. D. Hollenbach & H. Thronson (Dordrecht: Reidel), 171
- Heiles, C. 1996, in *Polarimetry of the Interstellar Medium*, ed. W. G. Roberge & D. C. B. Whittet (San Francisco: ASP), 457
- Heiles, C. 2000, *AJ*, 119, 923
- Hiltner, W. A. 1956, *ApJS*, 2, 389
- Höflich, P. 1991, *A&A*, 246, 481
- Höflich, P., Wheeler, J. C., Hines, D. C., & Trammell, S. R. 1996, *ApJ*, 459, 307
- Hough, J. H. 1996, in *Polarimetry of the Interstellar Medium*, ed. W. G. Roberge & D. C. B. Whittet (San Francisco: ASP), 569
- Jeffery, D. J. 1991a, *ApJ*, 375, 264
- Jeffery, D. J. 1991b, *ApJS*, 77, 405
- Johnson, P. E. 1982, *Nature*, 295, 371
- Jones, T. J. 1989, *AJ*, 98, 2062
- Jones, T. J. 1996, in *Polarimetry of the Interstellar Medium*, ed. W. G. Roberge & D. C. B. Whittet (San Francisco: ASP), 381
- Jones, T. J., Klebe, D., & Dickey, J. M. 1992, *ApJ*, 389, 602
- Kartje, J. F. 1995, *ApJ*, 452, 565
- Kawabata, K. S., et al. 2002, *ApJ*, submitted (astro-ph/0205414)
- Leonard, D. C., & Filippenko, A. V. 2001, *PASP*, 113, 920 (LF01)
- Leonard, D. C., Filippenko, A. V., Ardila, D. R., & Brotherton, M. S. 2001, *ApJ*, 553, 861
- Leonard, D. C., Filippenko, A. V., Barth, A. J., & Matheson, T. 2000a, *ApJ*, 536, 239
- Leonard, D. C., Filippenko, A. V., Chornock, R., & Foley, R. J. 2002a, *PASP*, submitted (astro-ph/0206369)
- Leonard, D. C., et al. 2002b, *PASP*, 114, 35
- Leonard, D. C., et al. 2002c, *AJ*, in press (L02) (astro-ph/0207601)

- Leonard, D. C., Filippenko, A. V., & Matheson, T. 2000b, in *Cosmic Explosions*, ed. S. S. Holt & W. W. Zhang (New York: AIP), 165
- Mathewson, D. S., & Ford, V. L. 1970, *MmRAS*, 74, 139
- Mathis, J. S. 1986, *ApJ*, 308, 281
- McCall, M. L. 1984, *MNRAS*, 210, 829
- Miller, J. S., & Goodrich, R. W. 1990, *ApJ*, 355, 456
- Miller, J. S., Goodrich, R. W., & Mathews, W. G. 1991, *ApJ*, 378, 47
- Miller, J. S., & Stone, R. P. S. 1993, *Lick Obs. Tech. Rep.*, No. 66
- Oke, J. B., et al. 1995, *PASP*, 107, 375
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical Recipes in C: The Art of Scientific Computing*, 2nd edition (Cambridge: Cambridge University Press)
- Rank, D. M., Bregman, J., Witteborn, F. C., Cohen, M., Lynch, D. K., & Russell, R. W. 1988, *ApJ*, 325, L1
- Rodrigues, C. V., Magalhães, A. M., Coyne, G. V., & Piirola, V. 1997, *ApJ*, 485, 618
- Savage, B. D., & Mathis, J. S. 1979, *ARA&A*, 17, 73
- Scarrott, S. M., Rolph, C. D., & Semple, D. P. 1990, in *Galactic and Intergalactic Magnetic Fields*, ed. R. Beck, P. P. Kronberg, & R. Wielebinski (Dordrecht: Kluwer), 245
- Scarrott, S. M., Rolph, C. D., Wolstencroft, R. W., & Tadhunter, C. N. 1991, *MNRAS*, 249, 16
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schmidt, G. D., Elston, R., & Lupie, O. L. 1992, *AJ*, 104, 1563
- Serkowski, K. 1973, in *Interstellar Dust and Related Topics*, ed. J. M. Greenberg and H. C. van de Hulst (Dordrecht: Reidel), 115
- Serkowski, K., Mathewson, D. S., & Ford, V. L. 1975, *ApJ*, 196, 261
- Spitzer, L., Jr. 1978, *Physical Processes in the Interstellar Medium* (New York: John Wiley & Sons)

- Sukumar, S., & Allen, R. J. 1991, *ApJ*, 382, 100
- Teerikorpi, P. 1990, *A&A*, 235, 362
- Trammell, S. R., Hines, D. C., & Wheeler, J. C. 1993, *ApJ*, 414, 21
- Tran, H. D. 1995, *ApJ*, 440, 565
- Tran, H. D., Filippenko, A. V., Schmidt, G. D., Bjorkman, K. S., Jannuzi, B. T., & Smith, P. S. 1997, *PASP*, 109, 489
- Wang, L., & Wheeler, J. C. 1996, *ApJ*, 462, L27
- Wheeler, J. C. 2000, in *Cosmic Explosions*, ed. S. S. Holt & W. W. Zhang (New York: AIP), 445
- Whittet, D. C. B. 1992, *Dust in the Galactic Environment* (Philadelphia: Institute of Physics Publishing, Inc.)
- Whittet, D. C. B., Martin, P. G., Hough, J. H., Rouse, M. F., Bailey, J. A., & Axon, D. J. 1992, *ApJ*, 386, 562
- Whittet, D. C. B., & van Breda, I. G. 1978, *A&A*, 66, 57
- Wiling, B. A., Lebofsky, M. J., & Rieke, G. H. 1982, *AJ*, 87, 695
- Wooden, D. H. 1997, in *Astrophysical Implications of the Laboratory Study of Presolar Materials*, ed. T. J. Bernatowicz & E. Zinner (New York: AIP), 317
- Wooden, D. H., Rank, D. M., Bregman, J. D., Witteborn, F. C., Tielens, A. G. G. M., Cohen, M., Pinto, P. A., & Axelrod, T. S. 1993, *ApJS*, 88, 477